

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/328808926>

THE EFFECTS OF POST-ACTIVATION POTENTIATION ON UPPER-BODY POWER PERFORMANCE IN RECREATIONALLY TRAINED MEN

Article in *Homosporticus* · June 2018

CITATIONS

0

READS

106

5 authors, including:



Mensur Vrcić

University of Sarajevo

52 PUBLICATIONS 15 CITATIONS

[SEE PROFILE](#)



Erol Kovačević

University of Sarajevo

55 PUBLICATIONS 14 CITATIONS

[SEE PROFILE](#)



Denis Čaušević

University of Sarajevo

25 PUBLICATIONS 13 CITATIONS

[SEE PROFILE](#)



Amila Hodžić

University of Sarajevo

4 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Fitness-individualni programi [View project](#)



Second South – East Europe Conference of Martial Arts and Combat Sports - „ESP 2017 - Education, Science and Practice“ [View project](#)



HOMO SCIENTIFIC JOURNAL OF SPORT AND PHYSICAL EDUCATION **SPORTICUS**

UDC 796.011 **VOLUME 20 - ISSUE 1 - JUNE, 2018.**

ISSN 1512-8822



fasto

FACULTY OF SPORT AND
PHYSICAL EDUCATION
UNIVERSITY OF SARAJEVO

THE EFFECTS OF POST-ACTIVATION POTENTIATION ON UPPER-BODY POWER PERFORMANCE IN RECREATIONALLY TRAINED MEN

¹ Faculty of Sport and Physical Education, University of Sarajevo, Bosnia and Herzegovina

² Institute of Sport, Faculty of Sport and Physical Education, University of Sarajevo, Bosnia and Herzegovina

³ Center for Education and Entrepreneur, Sarajevo, Bosnia and Herzegovina

Original scientific paper
UDC: 796.433.7.012.11

Abstract

The aim of this study was to determine the optimal rest period between PAP and medicine ball throwing, and to verify that the performance of the explosive strength of upper limbs was improved when it was preceded by 3x3 90% of 1RM on the incline bench press. Fifteen male respondents (age = 21 ± 1.53 ; body mass = 73.75 ± 5.18) students of Faculty of Sport and Physical Education, University of Sarajevo, participated in this research. Respondents throw the ball from sitting position, and 3, 5, 7, 10 minutes after the activation stimulus - incline bench press 3x3 90% 1RM. Analyzing the obtained results, an acute increase in the distance of the medicine ball throw was detected. The highest statistically significant pre and post-AP changes occurred in the seventh minute and amounted to 17 cm and 2.89% ($p = 0.033$). 3x3-90% 1RM IBP results in statistically significant PAP effects and optimal recovery time is 7 minutes.

Key words: strength, students, PAP, medicine ball, 1RM

Introduction

Improving new training strategies, in the focus as a primary aim sets the promotion of the strong performance of athletes. One of the most widely used and most applicable methods of improving these performances is PAP (Post-activation Potentiation) and implies an acute increase in the effectiveness of muscular contraction after previous muscular activity (Ebben, W. P., 2006; do Carmo et al. 2018). Just increasing this efficiency accompanied by neuromuscular changes can contribute improving performance in exercises that require strength, speed or power (Borba et al. 2017; MacIntosh et al. 2012). However, various studies revealed that PAP does not appear equally in all athletes, and that there are different parameters influencing the size of its effect, eg gender (Jensen and Ebben 2003; Rixon et al. 2007), training level (Wilson et al. 2013; Chiu et al. 2003, Pääsuke et al. 2007), muscular fiber (Hamada et al. 2000; Seitz et al. 2016), duration of contraction (Baudry and Duchateau 2004; Shima et al. 2006) etc. Also, the individual characteristics of athletes such as level of training, distribution and the ratio of different types of muscle fibers affect the size of the PAP effect (Evetovich et al., 2015).

In today's research it is not clearly defined stacking size that influences the strength of the occurrence and size of the PAP effect but the correlation, which is confirmed in many studies (Duthie et al. 2002; Kilduff et al. 2007; Xenofontos et al. 2018; Lockie et al. 2018). The study Bellar et al. (2012) noticed a positive correlation between absolute power and the distance of throwing the ball after activating stimulus by throwing balls of bigger weight. Jo et al. (2010) state that absolute strength has an impact on time course of PAP effect, and that stronger people need less time to complete recovery. Also, Seitz et al. (2014) noticed the greatest PAP effect occurs 3 minutes after activating stimulus for stronger (1RM squat $\geq 2 \times$ body mass) and 6 minutes for weaker subjects. However, study shows that there is no precise optimum time of PAP effect, especially when it comes to the power of the upper extremities. There are indicators that the best effects are achieved between 7 minutes (de Assis Ferreira et al., 2012; Bevan et al. 2009) and 12 minutes (Kilduff et al. 2007). To be more precise, when it comes to acute improvements in ballistic movement performance, some authors report the statis-

tically significant improvements (Markovic et al., 2008; French et al., 2003; Smilios et al., 2005), while other authors have not noticed any improvement (Scott and Docherty, 2004; Jones and Lee, 2003; Koch et al., 2003). Therefore, the general aim of this study was to verify that the performance of the explosive strength of the upper limbs was improved when it was preceded by 3x3 90% of the 1RM on incline bench press. The second purpose was to determine the optimal rest period between PAP and medicine ball throwing.

Methods

Participants

The sample subjects of this study consisted of 15 healthy, physically active male students of Faculty of Sport and Physical Education of the University of Sarajevo, without any reported upper extremity injuries in the past two years. The subjects had experience with strength training and were all physically active and had 3 to 5 trainings per week. The characteristics of subjects (mean $\pm$ SD) were: age (21 ± 1.53), body mass (73.75 ± 5.18), maximum incline bench press 1RM strength - (66 ± 10.38). All subjects were previously informed in detail about the reasons and procedures of research and they signed a written consent, voluntarily.

Experimental Approach to the Problem

The subjects were asked not to do heavy training at least 24 hours before participating in this study. The research was conducted in two working days in the morning, between with one-day break. The first morning measured the weight of the examinees (BC-420MA, TANITA Europe GmbH, Sindelfingen, Germany), after which they performed a standard warm-up of 5 minutes of easy running ($8 \text{ km} \cdot \text{h}^{-1}$) followed by 5 minutes of dynamic stretching (Tsoukos et al. 2016) after which 1RM was determined.

One-Repetition Maximum Bench Press

Determining maximum dynamic strength (1RM) was made according to the recommendation of Brown and Weir (2001) with the help of barbell and free weights. The subjects were asked to lie down on the incline press (28° incline) with head, shoulder blades, and buttocks constantly contacting the bench. The bar is lowered to the chest parallel to the nipple line, after which one pushes upward until the elbows aren't fully corrected. The 1RM was determined by using the standard procedure (Brown and Weir, 2001; Dohoney et al., 2002). After determining 90% of 1RM, the loads are gradually increased with the dumbbells of 1.25; 2.5 and 5 kg until 1RM is reached. For all subjects 90% of 1RM corresponded to their 3RM.

Seated Medicine Ball Throw

The second day included warm-up and throwing the medicine ball (4kg). The medicine ball throwing test used is recommended by the authors Harasin et al. 2006, which was

performed by the subject sitting on a chair with a medicine ball (4 kg, medical volume 76 cm) in both hands on the chest. The subject was asked to throw the medicine as far as possible in a certain direction and after then the distance of the throw have been measured. There were two throws in 10 second intervals in order to avoid the acute effects of the first throwing on the second. This test is used because it is very simple and highly reproducible test of upper-body explosive performance (int. Corell = 0.97) (Harasin et al. 2006), and very often used in the training of athletes in many sports. Activation stimulus meant 3x3 with 90% 1RM after which they threw the medicine ball at certain time points of 3, 5, 7 and 10 minutes. Results of throwing distance medicine ball are marked and rounded to the lower whole number.

Statistical Analysis

The results were processed using SPSS. 23 for Windows (IBM Corp. Chicago). Arithmetic mean values and standard deviations were calculated, for all variables, followed by paired samples T-test to determine possible differences in time points. Statistical significance deemed at $p < 0.05$.

Results


Table 1 shows the results of arithmetic means and standard deviations of treated variables. It is noticeable that the results of the pair sample T-test show statistically significant difference in the power output by throwing medicine ball only in the seventh minute ($p = .033$). Figure 1. shows the average means of the distance of medical ball throwing at certain points that were analyzed in this study.

Table 1 Descriptive values of the achieved results ($n = 15$)

Variables	Mean	Std. Deviation	T - test
Weight	73.75	5.188	/
IBP	66.00	10.385	/
PRE - PAP	5.89	.4760	/
MBP 3	5.97	.376	.380
MBP 5	6.04	.353	.109
MBP 7	6.06	.444	.033*
MBP 10	5.99	.452	.182

* significant at $p < 0.05$; IBP = Incline bench press; PRE - PAP = medicine ball distance before PAP; MBP3 = distance after 3 minutes; MBP5 = distance after 5 minute; MBP7 = distance after 7 minute; MBP10 = distance after 10 minute;

Figure 1 Mean values of the distance of medical ball throwing (3,5,7,10min)



Discussion

The aim of this study was to evaluate PAP effects during upper limb power exercise after a 3x3-90%1RM IBP. Two evaluated factors were (1) PAP magnitude and (2) optimal recovery time.

The most obvious acute effect of previous highly intensive or high-performing activities is the loss of the ability to generate force. However, it has been observed that contractile history can also result in improved performance. High intensity and/or high-volume physical activities primarily reduces force generation ability (MacIntosh i Rassier, 2002). However, it is noticed that this type of contractile history can also enhance performance for a short time (Smith i Fry, 2007; Grange et al, 1993; Sweeney et al, 1993). Previous studies have shown positive PAP effects on nerve activation magnitude (Hamada i sur., 2000; Moore i Stull, 1984; Suzuki i sur., 1988) and contractile characteristics (Hamada i sur., 2000; Vandervoort et al, 1983).

If observed in percentages, if the contraction (or contractile parameters) after the activation stimulus is 100% of the initial means, PAP does not occur, while means bigger than 100% indicate the presence of PAP and less than 100% indicates "posterior activation depression" (Mola et al., 2014) or fatigue domination over PAP effects (Chiu et al., 2003). Further in the text, the changes in measurements after the activation stimuli will be represented in percentages, compared to the initial measurement.

According to the aim and based on the results presented in the previous chapter, it is noticed that 3x3 90% 1RM incline bench press increased upper-body ballistic performance. The highest statistically significant pre and post-AP changes observed were 17 cm (2.89%; $p = .033$) seven minutes after AP while other changes, although present, were also positive but not statistically significant. Thus, the increase in medicine ball throw distance after three minutes was 9 cm (1.36%), 15 cm (2.55%) after five minutes and 10cm (1.70%) after 10 minutes.

These results are in line with the review article in which author (Bishop, 2003) who stated that medium (and stronger)

intensity activation stimuli may cause acute improvement in the performance of subsequent task performance until local muscular fatigue occurs. It is important to mention that Sale (2002) stated that PAP affects subsequent performance, but the optimal parameters are not fully determined.

In addition to review articles that point to the existence of PAP effects, Hrysomallis and Kidgell (2001) did not notice statistically significant difference in the scores recorded on the force plate during explosive push-ups after a 5RM bench press activation stimulus.

Liossis et al (2013) noted that strength output increases after 65% and 85% 1RM bench press, whereas at 65% 1RM changes occur after 4 minutes while at 85% 1RM the greatest PAP effects were observed 8 minutes after activation stimulation. These findings are consistent with the results of the study since, after 90% of 1RM, the biggest difference was observed 7 minutes after AS.

Considering that the factors affecting the level of PAP: 1) the age of the respondents (Baudry et al., 2005), 2) individual's training level (Chiu et al., 2003; Wilson et al., 2013), 3) the type of sport (Pääsuke i sur., 2007), 5) the ratio of fast and slow twitch muscle fibers (Hamada et al., 2000) and 6) contraction type (Ye et al., 2015) it is likely that the results would be a little different if sample with different characteristics was included in the study, for example professional athletes, athletes which predominantly use their upper extremities or participate in training involving mostly explosive activities.

Many argued about possible PAP mechanisms and concluded that this type of contraction is primarily associated with myosin regulatory light chain phosphorylation (Grange et al., 1993; Sweeney et al., 1993; Metzger et al., 1989) and an increase in α -motoneuron excitability (Misiasek, 2003, Zucker and Regehr, 2002), and, additionally, the increase in central nervous system stimulation (Rixon et al., 2007) which manifests itself by increasing nerve conduction and decreasing presynaptic inhibition thus resulting in better synchronization and activation of higher order motor units (Aagard et al., 2002; Chiu et al., 2003; Güllich and Schmidtbleicher, 1996; Tillin & Bishop, 2009)

In addition to the myosin regulatory light chain phosphorylation, some of the PAP mechanisms are increased motor-unit excitability, better motor-unit synchronization (Hodgson et al., 2005; Rassier and MacIntosh, 2000), acute changes in muscle fiber angle (Tillin and Bishop, 2009), increase in actin-myosin binding number and better interaction (Gülich and Schmidtbleicher, 1996), improved stimulation-contraction synchronization and increased motor-unit activation (Zhi et al., 2005; Szczena et al., 2002) Klein et al., 2001; de Luca et al., 1996).

Conclusion

In accordance with the set of aims, it can be concluded that 3x3-90% 1RM IBP results in a statistically significant increase in the distance of the medical ball throwing distance from the chest and the optimal recovery time after activation stimulus with the amateurs is 7 minutes.

References

- Aagaard, P., Simonsen, E. B., Andersen, J. L., Magnusson, P., & Dyhre-Poulsen, P. (2002). Neural adaptation to resistance training: changes in evoked V-wave and H-reflex responses. *Journal of Applied Physiology*, 92(6), 2309-2318.
- Baudry, S., & Duchateau, J. (2004). Postactivation potentiation in human muscle is not related to the type of maximal conditioning contraction. *Muscle & Nerve: Official Journal of the American Association of Electrodiagnostic Medicine*, 30(3), 328-336.
- Baudry, S., Klass, M., & Duchateau, J. (2005). Postactivation potentiation influences differently the nonlinear summation of contractions in young and elderly adults. *Journal of Applied Physiology*, 98(4), 1243-1250.
- Bellar, D., Judge, L. W., Turk, M., & Judge, M. (2012). Efficacy of potentiation of performance through overweight implement throws on male and female collegiate and elite weight throwers. *The Journal of Strength & Conditioning Research*, 26(6), 1469-1474.
- Bevan, H. R., Owen, N. J., Cunningham, D. J., Kingsley, M. I., & Kilduff, L. P. (2009). Complex training in professional rugby players: Influence of recovery time on upper-body power output. *The Journal of Strength & Conditioning Research*, 23(6), 1780-1785.
- Bishop, D. Warm up II: Performance changes following active warm up and how to structure the warm up. *Sports Med.* 33: 483-498. 2003.
- Borba, D. D. A., Ferreira-Júnior, J. B., Santos, L. A. D., Carmo, M. C. D., & Coelho, L. G. M. (2017). Effect of post-activation potentiation in Athletics: a systematic review. *Revista Brasileira de Cineantropometria & Desempenho Humano*, 19(1), 128-138.
- Brown, L. E., & Weir, J. P. (2001). ASEP procedures recommendation I: accurate assessment of muscular strength and power. *Journal of Exercise Physiology Online*, 4(3).
- Chiu, L. Z., Fry, A. C., Weiss, L. W., Schilling, B. K., Brown, L. E., & Smith, S. L. (2003). Postactivation potentiation response in athletic and recreationally trained individuals. *The Journal of Strength & Conditioning Research*, 17(4), 671-677.
- de Assis Ferreira, S. L., Panissa, V. L. G., Miarka, B., & Franchini, E. (2012). Postactivation potentiation: Effect of various recovery intervals on bench press power performance. *The Journal of Strength & Conditioning Research*, 26(3), 739-744.
- de Luca, C. J., Foley, P. J., & Erim, Z. (1996). Motor unit control properties in constant-force isometric contractions. *Journal of Neurophysiology*, 76(3), 1503-1516.
- do Carmo, E. C., De Souza, E. O., Roschel, H., Kobal, R., Ramos, H., Gil, S., ... & do Carmo, E. C. (2018). Self-Selected Rest Interval Improves Vertical Jump Post-Activation Potentiation. *The Journal of Strength & Conditioning Research*.
- Dohoney, P., Chromiak, J., Lemire, D., Abadie, B., & Kovacs, C. (2002). Prediction of one repetition maximum (1-RM) strength from a 4-6 RM and a 7-10 RM submaximal strength test in healthy young adult males. *J Exerc Physiol*, 5(3), 54-9.
- Duthie, G. M., Young, W. B., & Aitken, D. A. (2002). The acute effects of heavy loads on jump squat performance: an evaluation of the complex and contrast methods of power development. *The Journal of Strength & Conditioning Research*, 16(4), 530-538.
- Ebben, W. P. (2006). A brief review of concurrent activation potentiation: theoretical and practical constructs. *Journal of strength and Conditioning Research*, 20(4), 985.
- Evetovich, T. K., Conley, D. S., & McCawley, P. F. (2015). Postactivation potentiation enhances upper-and lower-body athletic performance in collegiate male and female athletes. *The Journal of Strength & Conditioning Research*, 29(2), 336-342.
- French, D. N., Kraemer, W. J., & Cooke, C. B. (2003). Changes in dynamic exercise performance following a sequence of preconditioning isometric muscle actions. *The Journal of Strength & Conditioning Research*, 17(4), 678-685.
- Grange, R. W., Vandenboom, R., & Houston, M. E. (1993). Physiological significance of myosin phosphorylation in skeletal muscle. *Canadian Journal of Applied Physiology*, 18(3), 229-242.
- Gülich, A., & Schmidtbleicher, D. (1996). MVC-induced short-term potentiation of explosive force. *New Studies in Athletics*, 11, 67-84.
- Hamada, T., Sale, D. G., MacDougall, J. D., & Tarnopolsky, M. A. (2000). Postactivation potentiation, fiber type, and

- twitch contraction time in human knee extensor muscles. *Journal of Applied Physiology*, 88(6), 2131-2137.
- Harasin, D., Dizdar, D., & Marković, G. (2006). High reliability of tests of maximum throwing performance. *Journal of Human Movement Studies*, 51(1), 63-76.
- Hodgson, M., Docherty, D., & Robbins, D. (2005). Post-activation potentiation. *Sports Medicine*, 35(7), 585-595.
- Hrysomallis, C., & Kidgell, D. (2001). Effect of heavy dynamic resistive exercise on acute upper-body power. *Journal of Strength and Conditioning Research*, 15(4), 426-430.
- Jensen, R. L., & Ebben, W. P. (2003). Kinetic analysis of complex training rest interval effect on vertical jump performance. *The Journal of Strength & Conditioning Research*, 17(2), 345-349.
- Jo, E., Judelson, D. A., Brown, L. E., Coburn, J. W., & Dabbs, N. C. (2010). Influence of recovery duration after a potentiating stimulus on muscular power in recreationally trained individuals. *The Journal of Strength & Conditioning Research*, 24(2), 343-347.
- Jones, P., & Lees, A. (2003). A biomechanical analysis of the acute effects of complex training using lower limb exercises. *Journal of Strength and Conditioning Research*, 17(4), 694-700.
- Kilduff, L. P., Bevan, H. R., Kingsley, M. I., Owen, N. J., Bennett, M. A., Bunce, P. J., ... & Cunningham, D. J. (2007). Postactivation potentiation in professional rugby players: optimal recovery. *The Journal of Strength & Conditioning Research*, 21(4), 1134-1138.
- Klein, C., Rice, C., & Marsh, G. (2001). Normalized force, activation, and coactivation in the arm muscles of young and old men. *Journal of Applied Physiology*, 91(3), 1341-1349.
- Koch, A. J., O'bryant, H. S., Stone, M. E., Sanborn, K., Proulx, C., Hruby, J., ... & Stone, M. H. (2003). Effect of warm-up on the standing broad jump in trained and untrained men and women. *The Journal of Strength & Conditioning Research*, 17(4), 710-714.
- Liossis, L. D., Forsyth, J., Liossis, C., & Tsolakis, C. (2013). The acute effect of upper-body complex training on power output of martial art athletes as measured by the bench press throw exercise. *Journal of human kinetics*, 39(1), 167-175.
- Lockie, R. G., Lazar, A., Davis, D. L., & Moreno, M. R. (2018). Effects of postactivation potentiation on linear and change-of-direction speed: analysis of the current literature and applications for the strength and conditioning coach. *Strength & Conditioning Journal*, 40(1), 75-91.
- MacIntosh, B. R., & Rassier, D. E. (2002). What is fatigue? *Canadian Journal of Applied Physiology*, 27(1), 42-55.
- MacIntosh, B. R., Robillard, M. E., & Tomaras, E. K. (2012). Should postactivation potentiation be the goal of your warm-up?. *Applied physiology, nutrition, and metabolism*, 37(3), 546-550.
- Markovic, G., Simek, S., & Bradic, A. (2008). Are acute effects of maximal dynamic contractions on upper-body ballistic performance load specific?. *The Journal of Strength & Conditioning Research*, 22(6), 1811-1815.
- Metzger, J. M., Greaser, M. L., & Moss, R. L. (1989). Variations in cross-bridge attachment rate and tension with phosphorylation of myosin in mammalian skinned skeletal muscle fibers. Implications for twitch potentiation in intact muscle. *The Journal of General Physiology*, 93(5), 855-883.
- Misiaszek, J. E. (2003). The H-reflex as a tool in neurophysiology: Its limitations and uses in understanding nervous system function. *Muscle & Nerve*, 28(2), 144-160.
- Moore, R. L., & Stull, J. T. (1984). Myosin light chain phosphorylation in fast and slow skeletal muscles in situ. *American Journal of Physiology-Cell Physiology*, 247(5), C462-C471.
- Pääsuke, M., Saapar, L., Ereline, J., Gapeyeva, H., Requena, B., & Ööpik, V. (2007). Postactivation potentiation of knee extensor muscles in power-and endurance-trained, and untrained women. *European Journal of Applied Physiology*, 101(5), 577-585.
- Rassier, D., & Macintosh, B. (2000). Coexistence of potentiation and fatigue in skeletal muscle. *Brazilian Journal of Medical and Biological Research*, 33(5), 499-508.
- Rixon, K. P., Lamont, H. S., & Bemben, M. G. (2007). Influence of type of muscle contraction, gender, and lifting experience on postactivation potentiation performance. *The Journal of Strength & Conditioning Research*, 21(2), 500-505.
- Sale, D.G. Postactivation potentiation: Role in human performance. *Exerc. Sport Sci. Rev.* 30:138–143. 2002
- Scott, S. L., & Docherty, D. (2004). Acute effects of heavy preloading on vertical and horizontal jump performance. *Journal of Strength and Conditioning Research*, 18(2), 201-205.
- Seitz, L. B., de Villarreal, E. S., & Haff, G. G. (2014). The temporal profile of postactivation potentiation is related to strength level. *The Journal of Strength & Conditioning Research*, 28(3), 706-715.
- Seitz, L. B., Trajano, G. S., Haff, G. G., Dumke, C. C., Tufano, J. J., & Blazevich, A. J. (2016). Relationships between maximal strength, muscle size, and myosin heavy chain isoform composition and postactivation potentiation. *Applied Physiology, Nutrition, and Metabolism*, 41(5), 491-497.
- Shima, N., Rice, C. L., Ota, Y., & Yabe, K. (2006). The effect of postactivation potentiation on the mechanomyogram. *European journal of applied physiology*, 96(1), 17-23.

- Smilios, I., Piliandis, T., Sotiropoulos, K., Antonakis, M., & Tokmakidis, S. P. (2005). Short-term effects of selected exercise and load in contrast training on vertical jump performance. *The Journal of Strength & Conditioning Research*, 19(1), 135-139.
- Smith, J. C., & Fry, A. C. (2007). Effects of a ten-second maximum voluntary contraction on regulatory myosin light-chain phosphorylation and dynamic performance measures. *The Journal of Strength & Conditioning Research*, 21(1), 73-76.
- Suzuki, S., Kaiya, K., Watanabe, S., & Hutton, R. S. (1988). Contraction-induced potentiation of human motor unit discharge and surface EMG activity. *Medicine and Science in Sports and Exercise*, 20(4), 391-395.
- Sweeney, H., Bowman, B. F., & Stull, J. T. (1993). Myosin light chain phosphorylation in vertebrate striated muscle: regulation and function. *American Journal of Physiology-Cell Physiology*, 264(5), C1085-C1095.
- Szczesna, D., Zhao, J., Jones, M., Zhi, G., Stull, J., & Potter, J. D. (2002). Phosphorylation of the regulatory light chains of myosin affects Ca^{2+} sensitivity of skeletal muscle contraction. *Journal of Applied Physiology*, 92(4), 1661-1670.
- Tillin, N. A., & Bishop, D. (2009). Factors modulating post-activation potentiation and its effect on performance of subsequent explosive activities. *Sports Medicine*, 39(2), 147-166.
- Tsoukos, A., Bogdanis, G. C., Terzis, G., & Veligekas, P. (2016). Acute improvement of vertical jump performance after isometric squats depends on knee angle and vertical jumping ability. *Journal of strength and conditioning research*, 30(8), 2250-2257.
- Vandervoort, A., Quinlan, J., & McComas, A. (1983). Twitch potentiation after voluntary contraction. *Experimental Neurology*, 81(1), 141-152.
- Wilson, J. M., Duncan, N. M., Marin, P. J., Brown, L. E., Loenneke, J. P., Wilson, S. M., . . . Ugrinowitsch, C. (2013). Meta-analysis of postactivation potentiation and power: effects of conditioning activity, volume, gender, rest periods, and training status. *The Journal of Strength & Conditioning Research*, 27(3), 854-859.
- Xenofondos, A., Bassa, E., Vrabas, I. S., Kotzamanidis, C., & Patikas, D. A. (2018). Muscle Twitch Torque During Two Different in Volume Isometric Exercise Protocols: Fatigue Effects on Postactivation Potentiation. *The Journal of Strength & Conditioning Research*, 32(2), 578-586.
- Ye, X., Beck, T. W., & Wages, N. P. (2015). Acute effects of concentric vs. eccentric exercise on force steadiness and electromyographic responses of the forearm flexors. *The Journal of Strength & Conditioning Research*, 29(3), 604-611.
- Zhi, G., Ryder, J. W., Huang, J., Ding, P., Chen, Y., Zhao, Y., . . . Stull, J. T. (2005). Myosin light chain kinase and myosin phosphorylation effect frequency-dependent potentiation of skeletal muscle contraction. *Proceedings of the National Academy of Sciences of the United States of America*, 102(48), 17519-17524.
- Zucker, R. S., & Regehr, W. G. (2002). Short-term synaptic plasticity. *Annual Review of Physiology*, 64(1), 355-405.

Submitted: April 11, 2018

Accepted: May 19, 2018

Corresponding author:

Mensur Vrcić, PhD

Faculty of sport and physical education,

University of Sarajevo

E-mail: mvrčić@fasto.unsa.ba